

# 1 Smoothness of $\text{Bun}_n$ for dinguses<sup>1</sup>

Dear Jared,

Let  $\text{Bun}_n \rightarrow \text{Perf}$  denote the stack of rank  $n$  vector bundles on “the” Fargues-Fontaine curve. Yesterday I figured out a fairly cheap argument for checking that  $\text{Bun}_n$  is a smooth diamond stack, using charts made out of de Rham affine Grassmannians. Of course Peter’s charts made from those spaces  $X_b$  give more information, but it seems harder to check that they have the right properties.

A word on terminology: if  $S$  is any absolute diamond, we say  $S$  is *smooth* if for any diamond  $X$ , the projection map  $X \times S \rightarrow X$  is smooth. Note that if  $S_1$  and  $S_2$  are smooth, then so is  $S_1 \times S_2$ . One can also check that if  $K$  is any finite extension of  $\mathbf{Q}_p$  and  $S$  is a diamond with a smooth morphism  $S \rightarrow \text{Spd } K$ , then  $S$  is smooth in this sense.

Let  $\text{Bun}_n^d \subset \text{Bun}_n$  denote the open-closed substack of bundles of constant degree  $d$ . Let  $\text{Gr}_{n,k}/\text{Spd } \mathbf{Q}_p$  denote the de Rham affine Grassmannian sending  $T \in \text{Perf}$  with specified untilt  $T^\sharp$  to the set of subsheaves

$$\mathcal{E} \subset \mathcal{O}_{\mathcal{X}_T}^n$$

such that  $\mathcal{E} \rightarrow \mathcal{O}_{\mathcal{X}_T}^n$  is a modification supported along  $T^\sharp \subset \mathcal{X}_T$  of (constant) meromorphy type  $(k, 0, \dots, 0)$ . Note that  $\mathcal{E}$  has constant degree  $-k$ . In particular, for any  $m \geq d/n$  there is a natural morphism

$$\text{Gr}_{n,mn-d} \rightarrow \text{Bun}_{n,d}$$

given by sending  $\mathcal{E} \subset \mathcal{O}_{\mathcal{X}_T}^n$  as above to the degree  $d$  bundle  $\mathcal{E}(m) := \mathcal{E} \otimes_{\mathcal{O}} \mathcal{O}(m)$ . This clearly factors through a morphism

$$f_m : [\text{Gr}_{n,mn-d}/\underline{\text{GL}}_n(\mathbf{Q}_p)] \rightarrow \text{Bun}_{n,d},$$

where  $\underline{\text{GL}}_n(\mathbf{Q}_p)$  acts on any  $\text{Gr}_{n,k}$  in the usual way.

**Proposition 1.1.** *The morphism  $f_m$  is smooth.*

*Proof.* We need to check that for any  $S \in \text{Perf}$  and any morphism  $a : S \rightarrow \text{Bun}_{n,d}$ , the fiber product

$$S \times_{a, \text{Bun}_{n,d}, f_m} [\text{Gr}_{n,mn-d}/\underline{\text{GL}}_n(\mathbf{Q}_p)]$$

is a diamond smooth over  $S$ . What functor does this fiber product represent? Well, giving  $a$  is equivalent to giving a degree  $d$  rank  $n$  bundle  $\mathcal{E}/\mathcal{X}_S$ . Unwinding definitions then shows that this fiber product represents the set of isomorphism classes of pairs  $(S^\sharp, \mathcal{E} \hookrightarrow \mathcal{F})$  where  $S^\sharp$  is an untilt of  $S$  and  $\mathcal{E} \hookrightarrow \mathcal{F}$  is a modification supported along  $S^\sharp \subset \mathcal{X}_S$  and of meromorphy type  $(mn-d, 0, \dots, 0)$ , such that moreover  $\mathcal{F}$  is pointwise-semistable.<sup>2</sup> Ignoring the last condition, this functor is representable by a “twisted de Rham affine Grassmannian”  $\text{Gr}_{n,d-mn}^\mathcal{E}/S$ , which locally on  $S$  is isomorphic to  $\text{Gr}_{n,d-mn} \times S$  and therefore is smooth over  $S$ . Enforcing the semistability of  $\mathcal{F}$  then cuts out (by Kedlaya-Liu) an open subspace

$$\text{Gr}_{n,d-mn}^{\mathcal{E},ss} \subset \text{Gr}_{n,d-mn}^\mathcal{E},$$

so  $\text{Gr}_{n,d-mn}^{\mathcal{E},ss} \rightarrow S$  is still smooth, and

$$\text{Gr}_{n,d-mn}^{\mathcal{E},ss} \cong S \times_{\text{Bun}_{n,d}} [\text{Gr}_{n,mn-d}/\underline{\text{GL}}_n(\mathbf{Q}_p)]$$

so we win. □

---

<sup>1</sup>Version of 12/25/2016

<sup>2</sup>More precisely, this fiber product should be regarded as a functor on  $\text{Perf}/S$ , but whatever.

Next we describe the image of  $f_m$  on geometric points.

**Proposition 1.2.** *Let  $C/\mathbf{F}_p$  be an algebraically closed perfectoid field, and let  $a : \mathrm{Spd} C \rightarrow \mathrm{Bun}_{n,d}$  be any point, with associated bundle  $\mathcal{E}/\mathcal{X}_C$ . Then  $a$  lifts along  $f_m$  to a  $C$ -point of  $[\mathrm{Gr}_{n,mn-d}/\underline{\mathrm{GL}}_n(\mathbf{Q}_p)]$  if and only if the maximal Harder-Narasimhan slope of  $\mathcal{E}$  is  $\leq m$ .*

*Proof.* “Only if” is an easy exercise: if  $a$  lifts, then by definition there is some inclusion  $\mathcal{E}(-m) \subset \mathcal{O}_{\mathcal{X}_C}^n$ , so  $\mathcal{E}(-m)$  has maximal HN slope  $\leq 0$ . “If” can be deduced from various results of the form “weakly admissible filtrations of specified Hodge type on specified  $\varphi$ -modules exist when they should”.  $\square$

The condition on HN slopes in the previous proposition cuts out an open substack  $\mathrm{Bun}_{n,d}^{\leq m}$  such that  $f_m$  factors through the inclusion of this substack. Clearly  $\mathrm{Bun}_{n,d}^{\leq m} \subset \mathrm{Bun}_{n,d}^{\leq m+1}$  and

$$\mathrm{Bun}_{n,d} = \bigcup_{m \gg 0} \mathrm{Bun}_{n,d}^{\leq m}.$$

It is true, but not *a priori* obvious, that  $f_m : [\mathrm{Gr}_{n,mn-d}/\underline{\mathrm{GL}}_n(\mathbf{Q}_p)] \rightarrow \mathrm{Bun}_{n,d}^{\leq m}$  is surjective in the pro-étale topology, i.e. that given any  $S \in \mathrm{Perf}$  and any  $x \in \mathrm{Bun}_{n,d}^{\leq m}(S)$  we can lift  $x$  along  $f_m$  after passing to some pro-étale cover of  $S$ . This can be deduced as follows: Using the previous two proposition, one first checks that the morphism of diamonds

$$S \times_{s, \mathrm{Bun}_{n,d}^{\leq m}, f_m} [\mathrm{Gr}_{n,mn-d}/\underline{\mathrm{GL}}_n(\mathbf{Q}_p)] \rightarrow S$$

is smooth, and moreover surjective on topological spaces, with locally spatial source. One then applies the following result (whose straightforward proof is omitted; the key point in the proof is that smooth maps of diamonds are universally open).

**Proposition 1.3.** *Let  $f : Y \rightarrow X$  be any map of locally spatial diamonds. If  $f$  is smooth and  $|Y| \rightarrow |X|$  is surjective, then  $f$  is surjective as a map of pro-étale sheaves.*

OK, so we have a family of smooth maps

$$f_m : [\mathrm{Gr}_{n,mn-d}/\underline{\mathrm{GL}}_n(\mathbf{Q}_p)] \rightarrow \mathrm{Bun}_{n,d}$$

which together cover the target. Now comes the fun part.

**Proposition 1.4.** *The stack  $[\mathrm{Gr}_{n,mn-d}/\underline{\mathrm{GL}}_n(\mathbf{Q}_p)]$  is a smooth diamond stack.*

With this in hand, we’re done: after choosing some smooth diamonds  $X_m$  with some smooth surjective maps

$$g_m : X_m \rightarrow [\mathrm{Gr}_{n,mn-d}/\underline{\mathrm{GL}}_n(\mathbf{Q}_p)],$$

the composite maps  $f_m \circ g_m : X_m \rightarrow \mathrm{Bun}_{n,d}$  are smooth and give a collection of charts which verify that  $\mathrm{Bun}_{n,d}$  is a smooth diamond stack.

So now we need to show that  $[\mathrm{Gr}_{n,mn-d}/\underline{\mathrm{GL}}_n(\mathbf{Q}_p)]$  is smooth. We’d like to deduce this from the smoothness of  $\mathrm{Gr}_{n,k}$ . It turns out there’s a really cute general argument for this sort of thing (which is what I missed until yesterday).

**Proposition 1.5.** *Fix a locally profinite group  $G$ , and let  $X$  be any absolute diamond with  $\underline{G}$ -action. If there exists some smooth diamond  $W$  with a free  $\underline{G}$ -action, then  $[X/\underline{G}]$  is a diamond stack. If moreover  $W$  can be chosen such that  $W/\underline{G}$  is smooth, then  $[X/\underline{G}]$  is smooth whenever  $X$  is smooth.*

*Proof.* Give  $X \times W$  the diagonal  $\underline{G}$ -action; this action is free, since the action on  $W$  is free, so  $(X \times W)/\underline{G}$  is a diamond. The projection map  $X \times W \rightarrow X$  is smooth, surjective and  $\underline{G}$ -equivariant, so we get a smooth surjective map

$$(X \times W)/\underline{G} \rightarrow [X/\underline{G}]$$

whose source is a diamond.<sup>3</sup> Hence the target is a diamond stack.<sup>4</sup>

Suppose now that  $X$  is smooth. The natural projection map  $(X \times W)/\underline{G} \rightarrow W/\underline{G}$  is then smooth. Indeed, we get a pullback diagram

$$\begin{array}{ccc} X \times W & \longrightarrow & W \\ \downarrow & & \downarrow \\ (X \times W)/\underline{G} & \longrightarrow & W/\underline{G} \end{array}$$

with surjective pro-étale vertical maps, and smoothness of  $X$  implies that the upper horizontal map is smooth; since smoothness can be checked (quasi-)pro-étale-locally on the target, we get that the lower horizontal map is smooth as desired. But now, if  $W/\underline{G}$  is smooth as well, we're looking at a smooth map  $(X \times W)/\underline{G} \rightarrow W/\underline{G}$  with smooth target, which implies that  $(X \times W)/\underline{G}$  is smooth. But then  $(X \times W)/\underline{G} \rightarrow [X/\underline{G}]$  is a smooth surjective map whose source is a smooth diamond, so we win.  $\square$

Returning to our specific situation, we just need to find *some* smooth diamond  $W$  with a free  $\underline{\mathrm{GL}}_n(\mathbf{Q}_p)$ -action, such that  $W/\underline{\mathrm{GL}}_n(\mathbf{Q}_p)$  is also smooth. To do this, suppose we can find smooth diamonds  $W_1$  and  $W_2$ , where  $W_1$  has a free  $\underline{\mathrm{SL}}_n(\mathbf{Q}_p)$ -action and  $W_2$  has a free  $\mathbf{Q}_p^\times$ -action, such that  $W_1/\underline{\mathrm{SL}}_n(\mathbf{Q}_p)$  and  $W_2/\mathbf{Q}_p^\times$  are both smooth. Letting  $m : \underline{\mathrm{SL}}_n(\mathbf{Q}_p) \times \mathbf{Q}_p^\times \rightarrow \underline{\mathrm{GL}}_n(\mathbf{Q}_p)$  be the group homomorphism which is inclusion on the first factor and which sends  $(1, a)$  to  $\mathrm{diag}(a, \dots, a)$ , the diamond

$$W = (W_1 \times W_2) \times_{\underline{\mathrm{SL}}_n(\mathbf{Q}_p) \times \mathbf{Q}_p^\times} \underline{\mathrm{GL}}_n(\mathbf{Q}_p)$$

then does what we want: since  $\ker m$  is finite and  $\mathrm{im} m \subset \underline{\mathrm{GL}}_n(\mathbf{Q}_p)$  is a finite-index normal subgroup,  $W$  is étale over the smooth diamond  $W_1 \times W_2$ , hence smooth itself, and

$$W/\underline{\mathrm{GL}}_n(\mathbf{Q}_p) \cong W_1/\underline{\mathrm{SL}}_n(\mathbf{Q}_p) \times W_2/\mathbf{Q}_p^\times$$

is smooth.

For  $W_2$ , we just take  $\mathrm{Spd} \mathbf{Q}_p^{\mathrm{cyc}} \cong \mathrm{Spd} \mathbf{F}_p((t^{1/p^\infty}))$  with the usual  $\mathbf{Q}_p^\times$ -action. For  $W_1$ , it turns out that the following thing works. Let  $W_1$  be the functor on  $\mathrm{Perf}$  sending  $S$  to the set of pointwise-injective bundle maps  $i : \mathcal{O}^n \hookrightarrow \mathcal{O}(\frac{1}{n+1})$  over the relative curve  $\mathcal{X}_S$ . There is an obvious  $\underline{\mathrm{GL}}_n(\mathbf{Q}_p)$ -action given by precomposition with  $i$ . I claim that  $W_1$  and  $W_1/\underline{\mathrm{SL}}_n(\mathbf{Q}_p)$  are smooth.<sup>5</sup>

<sup>3</sup>This follows from a general lemma: If  $P$  is some property of morphisms of diamonds which is stable under base change and quasi-pro-étale-local on the target, and  $Y \rightarrow X$  is a  $\underline{G}$ -equivariant morphism of absolute diamonds which has  $P$ , then  $[Y/\underline{G}] \rightarrow [X/\underline{G}]$  has  $P$ , in the sense that for any diamond  $W$  with a map  $W \rightarrow [X/\underline{G}]$ ,  $[Y/\underline{G}] \times_{[X/\underline{G}]} W \rightarrow W$  has  $P$ .

<sup>4</sup>One also checks that  $[X/\underline{G}]$  always has diagonal representable in diamonds, for any absolute diamond with  $\underline{G}$ -action, cf. the “Notes on diamonds”.

<sup>5</sup>It seems very likely that  $W_1/\underline{\mathrm{GL}}_n(\mathbf{Q}_p)$  is actually smooth, in which case one could avoid the silly circumlocutions of the previous paragraph, but I wasn't able to see this smoothness immediately.

For the smoothness of  $W_1$ , consider the functor  $W'$  on Perf sending  $S$  to the set of sections  $s \in H^0(\mathcal{X}_S, \mathcal{O}(\frac{1}{n+1}))$  such that  $s$  does not vanish identically on any fiber of the map  $|\mathcal{X}_S| \rightarrow |S|$ . This functor is representable by a spatial diamond, which turns out by some games with Lubin-Tate formal modules to be of the shape  $\mathrm{Spd} \mathbf{F}_q((t^{1/p^\infty}))/\mathbf{Z}_{p^{n+1}}^\times$  for some free action of  $\mathbf{Z}_{p^{n+1}}^\times$  on some  $\mathrm{Spd} \mathbf{F}_q((t^{1/p^\infty}))$ ; in particular, this thing is smooth. (Here  $\mathbf{Z}_{p^h}$  = ring of integers in the degree  $h$  unramified extension of  $\mathbf{Q}_p$ .) Then  $W_1$  is an open subfunctor of  $\underbrace{W' \times \cdots \times W'}_n$ , so  $W_1$  is smooth.

For the smoothness of  $W_1/\mathrm{SL}_n(\mathbf{Q}_p)$ , we first observe that this thing has a moduli interpretation: it is the functor on Perf sending  $S$  to the set of pairs  $(\mathcal{E}, i)$  where  $\mathcal{E} \subset \mathcal{O}(\frac{1}{n+1})/\mathcal{X}_S$  is a rank  $n$  subbundle which is pointwise-semistable of degree zero and  $i$  is a trivialization  $i : \mathcal{O} \xrightarrow{\sim} \wedge^n \mathcal{E}$ . By some easy games with the classification, one can check that given any such  $\mathcal{E}$ ,  $\mathcal{O}(\frac{1}{n+1})/\mathcal{E}$  is a line bundle on  $\mathcal{X}_S$  of constant degree 1, and that  $i$  together with the trivialization  $\mathcal{O}(1) \cong \wedge^{n+1} \mathcal{O}(\frac{1}{n+1})$  induce a canonical trivialization  $\mathcal{O}(\frac{1}{n+1})/\mathcal{E} \cong \mathcal{O}(1)$ . Pushing this further,  $W_1/\mathrm{SL}_n(\mathbf{Q}_p)$  identifies with the functor sending  $S$  to the set of surjections  $\mathcal{O}(\frac{1}{n+1}) \twoheadrightarrow \mathcal{O}(1)$  of bundles over  $\mathcal{X}_S$ ; indeed, any such surjection has kernel  $\mathcal{E}$  which is pointwise-semistable of degree zero and which comes with a canonical trivialization of its determinant, and then  $W_1$  is the  $\mathrm{SL}_n(\mathbf{Q}_p)$ -torsor over this guy parametrizing trivializations  $\mathcal{O}^n \xrightarrow{\sim} \mathcal{E}$  compatible with the trivialization of  $\wedge^n \mathcal{E}$ . Applying  $\mathrm{Hom}_{\mathcal{O}_{\mathcal{X}_S}}(-, \mathcal{O}(1))$  to such a surjection gives an inclusion  $\mathcal{O} \hookrightarrow \mathcal{O}(\frac{n}{n+1})/\mathcal{X}_S$ , nonzero on each fiber of the map  $|\mathcal{X}_S| \rightarrow |S|$ , with cokernel  $\simeq \mathcal{O}(1)^n$  at all geometric points of  $S$ . In particular, we get a natural transformation

$$f : W_1/\mathrm{SL}_n(\mathbf{Q}_p) \rightarrow X$$

where  $X$  is the functor sending  $S$  to the set of sections  $s \in H^0(\mathcal{X}_S, \mathcal{O}(\frac{n}{n+1}))$  which are not identically zero on any fiber of  $|\mathcal{X}_S| \rightarrow |S|$ . I claim that  $f$  is an open immersion and that  $X$  is smooth. For openness, one easily checks that  $f$  is an injection. We then observe that  $f$  identifies its source with the subfunctor of its target cut out by the requirement that the vector bundle  $\mathcal{O}(\frac{n}{n+1})/\mathcal{O} \cdot s$  be pointwise-semistable, and the habitual openness of the latter condition gives what we want. Smoothness of  $X$ , finally, is analogous to the smoothness of  $W'$  and is left as an exercise.

Cheers,  
Dave